

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/342511049>

THE LONGHORNED BEETLES THAT ORIGINALLY DESCRIBED FROM WHOLE TERRITORIES OF TURKEY (COLEOPTERA: CERAMBYCOIDEA) PART II – CERAMBYCIDAE (CERAMBYCINAE AND STENOPTERINAE)

Article in *Munis Entomology and Zoology Journal* · June 2012

CITATIONS

4

READS

5

1 author:



[Hüseyin Özdikmen](#)

Gazi University

433 PUBLICATIONS 1,500 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Cerambycidae of Turkey [View project](#)



Chrysomelidae of Turkey [View project](#)

**THE LONGHORNED BEETLES THAT ORIGINALLY
DESCRIBED FROM WHOLE TERRITORIES OF TURKEY
(COLEOPTERA: CERAMBYCOIDEA) PART II –
CERAMBYCIDAE (CERAMBYCINAE AND STENOPTERINAE)**

Hüseyin Özdikmen*

* Gazi Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, 06500 Ankara, TURKEY. E-mail: ozdikmen@gazi.edu.tr

[Özdikmen, H. 2012. The longhorned beetles that originally described from whole territories of Turkey (Coleoptera: Cerambycoidea) Part II - Cerambycidae (Cerambycinae and Stenopterinae). *Munis Entomology & Zoology*, 7 (2): 699-707]

ABSTRACT. The paper is the second part of a planned serial work. It gives an annotated list of the Cerambycoidea [Cerambycidae (Cerambycinae and Stenopterinae)] that originally described from Turkey, with some important synonyms.

KEY WORDS: Cerambycinae, Stenopterinae, Cerambycidae, Cerambycoidea, Coleoptera, Turkey.

A series work is planned that is aim to expose the Turkish Cerambycoidea described from Turkey originally as possible as detailed and entirely by beginning from Vesperidae to Cerambycidae (Özdikmen, 2012a). The present study that includes the subfamilies Cerambycinae and Stenopterinae, is attempted as the second step of this aim.

Information in the present text is given in following order:

All taxa are accompanied by the author's name and description date. For each species, subspecies or important synonym, the original combination and type locality are provided under the taxon names. The status of endemism for Turkey is also indicated under these information.

In this paper, classification and nomenclature of the longhorn beetles suggested by Özdikmen (2012b) is followed chiefly. Within the genera and subgenera, the species are listed alphabetically.

Please see Löbl & Smetana (2010) for cited references in the text.

SUPERFAMILY CERAMBYCOIDEA Latreille, 1802

FAMILY CERAMBYCIDAE Latreille, 1802: 211

SUBFAMILY CERAMBYCINAE Latreille, 1802: 211

TRIBE HESPEROPHANINI Mulsant, 1839: 61

SUBTRIBE HESPEROPHANINA Mulsant, 1839: 61

GENUS *TRICHOFERUS* Wollaston, 1854: 427

SUBGENUS -

SPECIES *T. kotschyi* (Ganglbauer, 1883: 300)

Orig. comb.: *Hesperophanes kotschyi*

Type loc.: Asia Minor "Cilicia" (Turkey)

SPECIES *T. lunatus* (Szallies, 1994: 261)

Orig. comb.: *Hesperophanes lunatus*

Type loc.: Mardin: Hop pass (Turkey)

SPECIES *T. preissi* (Heyden, 1894: 85)

Orig. comb.: *Hesperophanes preissi*

Type loc.: Mardin (Turkey)

Endemic species to Turkey.

SPECIES *T. samai* Kadlec & Rejzek, 2001: 296

Orig. comb.: *Trichoferus samai*

Type loc.: Adiyaman: Nemrut Mt., Kahta env. (Turkey)

Endemic species to Turkey.

SPECIES *T. sbordonii* Sama, 1982: 217

Orig. comb.: *Trichoferus sbordonii*

Type loc.: Antalya: Alanya (Turkey)

Endemic species to Turkey.

TRIBE CERAMBYCINI Latreille, 1802: 211

SUBTRIBE CERAMBYCINA Latreille, 1802: 211

GENUS CERAMBYX Linnaeus, 1758: 388

SUBGENUS CERAMBYX Linnaeus, 1758: 388

SPECIES *C. heinzianus* Demelt, 1976: 65

Orig. comb.: *Cerambyx heinzianus*

Type loc.: Bingöl: Solhan (Turkey)

Endemic species to Turkey.

SUBGENUS MICROCERAMBYX Mikšić & Georgijevic, 1973: 22

SPECIES *C. scopolii* Fuessly, 1775: 12

Orig. comb.: *Cerambyx scopolii*

Type loc.: Switzerland

SUBSPECIES *C. s. nitidus* Pic, 1892: CXI

Orig. comb.: *Cerambyx scopolii* var. *nitidus*

Type loc.: "monts Amanus, pays d'Akbès" (Turkey: Hatay: Akbez)

Endemic subspecies to Turkey.

TRIBE ROSALIINI Fairmaire, 1864: 137

SUBTRIBE -

GENUS ROSALIA Audinet-Serville, 1834: 561

SUBGENUS ROSALIA Audinet-Serville, 1834: 561

SPECIES *R. alpina* (Linnaeus, 1758: 392)

Orig. comb.: *Cerambyx alpinus*

Type loc.: "Helvetia" (Switzerland)

SUBSPECIES *R. a. syriaca* Pic, 1895: CCLXXXV

Orig. comb.: *Rosalia alpina* var. *syriaca*

Type loc.: Hatay: Akbez (Turkey)

Endemic subspecies to Turkey.

TRIBE TRACHYDERINI Dupont, 1836: 1

SUBTRIBE TRACHIDERINA Dupont, 1836: 1

GENUS PURPURICENUS Dejean, 1821: 105

SUBGENUS -

SPECIES *P. bitlisiensis* Pic, 1902: 27

Orig. comb.: *Purpuricenus budensis* var. *bitlisiensis*

Type loc.: Bitlis (Turkey)

Endemic species to Turkey.

SPECIES *P. cornifrons* Sabbadini & Pesarini, 1992: 58

Orig. comb.: *Purpuricenus cornifrons*

Type loc.: Bingöl (Turkey)

Endemic species to Turkey.

SPECIES *P. interscapillatus* Plavilstshikov, 1937: 247 [RN]

Orig. comb.: *Purpuricenus interscapillatus*

Type loc.: Lebanon

SUBSPECIES *P. i. nudicollis* Demelt, 1968: 65

Orig. comb.: *Purpuricenus nudicollis*

Type loc.: İçel (Turkey)

Endemic subspecies to Turkey.

SPECIES *P. nigrotatatus* Pic, 1907: 169

Orig. comb.: *Purpuricenus budensis* var. *nigronotatus*

Type loc.: Adana: Karsanti (Turkey)

Endemic species to Turkey.

SYNONYM

konradi Bernhauer, 1976: 123

Orig. comb.: *Purpuricenus konradi*

Type loc.: Adana: Nurdağı pass (Turkey: Hatay prov.)

GENUS *CALCHAENESTHES* Kraatz, 1863: 97

SUBGENUS -

SPECIES *C. oblongomaculata* (Guerin-Meneville, 1844: 234)

Orig. comb.: *Callidium oblongomalatum*

Type loc.: Graecia (Greece)

SYNONYM

nogeli Frivaldszky, 1845: 183

Orig. comb.: *Callidium nogeli*

Type loc.: "Constantinopoly" (Turkey: İstanbul)

TRIBE CALLICHRMATINI Swainson & Shuckard, 1840: 293

SUBTRIBE -

GENUS *OSPHRANTERIA* Redtenbacher, 1850: 50

SUBGENUS -

SPECIES *O. coerulescens* Redtenbacher, 1850: 50

Orig. comb.: *Osphranteria coerulescens*

Type loc.: Shiraz (Iran)

SUBSPECIES *O. c. inaurata* Holzschuh, 1981: 98

Orig. comb.: *Osphranteria coerulescens inaurata*

Type loc.: Hakkari: Sat Mt.: Oramar (=Dağlıca) (Turkey)

TRIBE GRACILIINI Mulsant, 1839: 99

SUBTRIBE -

GENUS *HYBOMETOPIA* Ganglbauer, 1889: 282

SUBGENUS -

SPECIES *H. starcki* Ganglbauer, 1889: 285

Orig. comb.: *Hybometopia starcki*

Type loc.: Caucasus

SUBSPECIES *H. s. ivani* Sama, 1996: 106

Orig. comb.: *Hybometopia starcki ivani*

Type loc.: Kastamonu: Yarahgöz (Turkey)

TRIBE OBRINI Mulsant, 1839: 95

SUBTRIBE -

GENUS *ANATOLOBRIUM* Adlbauer, 2004: 419

SUBGENUS -

SPECIES *A. eggeri* Adlbauer, 2004: 421

Orig. comb.: *Anatolobrium eggeri*

Type loc.: Antalya: Alanya (Turkey)

Endemic species to Turkey.

TRIBE DELLINI Fairmaire, 1864: 154

SUBTRIBE -

GENUS *DELAGRANGEUS* Pic, 1892: XCIII

SUBGENUS *DELAGRANGEUS* Pic, 1892: XCIII

SPECIES *D. angustissimus* Pic, 1892: XCIII

Orig. comb.: *Delagrangaeus angustissimus*

Type loc.: "Haute Syria" (Turkey: Hatay: Akbez)

SYNONYM

asiaticus Pic, 1920: 21

Orig. comb.: *Mimophymatodes asiaticus*

Type loc.: Konya: Akşehir (Turkey)

SUBSPECIES *D. a. angustissimus* Pic, 1892: XCIII
Orig. comb.: *Delagrangeus angustissimus angustissimus*
Type loc.: "Haute Syria" (Turkey: Hatay: Akbez)
Endemic subspecies to Turkey.

GENUS *DEILUS* Audinet-Serville, 1834: 73

SUBGENUS -

SPECIES *D. rugosicollis* Rapuzzi & Sama, 2012: 668

Orig. comb.: *Deilus rugosicollis*

Type loc.: Lebanon, Syria, Turkey, Israel

TRIBE CALLIDIINI Kirby, 1837: 170

SUBTRIBE -

GENUS *ROPALOPUS* Mulsant, 1839: 40

SUBGENUS *ROPALOPUS* Mulsant, 1839: 40

SPECIES *R. hanae* Sama & Rejzek, 2002: 105

Orig. comb.: *Ropalopus hanae*

Type loc.: Muş: Buğlan pass (Turkey)

Endemic species to Turkey.

SPECIES *R. ledereri* (Fairmaire, 1866: 269)

Orig. comb.: *Rhopalopus ledereri*

Type loc.: Asie Mineure (Turkey: ?W Anatolia: ?İzmir)

SUBSPECIES *R. l. ledereri* (Fairmaire, 1866: 269)

Orig. comb.: *Rhopalopus ledereri*

Type loc.: Asie Mineure (Turkey: ?W Anatolia: ?İzmir)

SUBSPECIES *R. l. wittmeri* Demelt, 1970: 31

Orig. comb.: *Rhopalopus ledereri wittmeri*

Type loc.: Kahramanmaraş, Osmaniye (Turkey)

SPECIES *R. sculpturatus* (Pic, 1931: 9)

Orig. comb.: *Rhopalopus clavipes* var. *sculpturatus*

Type loc.: Tokat (Turkey)

Endemic species to Turkey.

GENUS *LEIODERES* Redtenbacher, 1849: 482

SUBGENUS -

SPECIES *L. tuerki* (Ganglbauer, 1886: 517)

Orig. comb.: *Callidium (Lioderes) türki*

Type loc.: Cilicia: Gülek (Bulgar-Dagh) (Turkey: İçel)

GENUS *CALLIDIUM* Fabricius, 1775: 187

SUBGENUS *CALLIDIUM* Fabricius, 1775: 187

SPECIES *C. syriacum* Pic, 1892: CXI

Orig. comb.: *Callidium ?aeneum* var. *syriacum*

Type loc.: "Nord Syrie" (Turkey: Hatay: Akbez not Syria)

GENUS *PHYMATODES* Mulsant, 1839: 47

SUBGENUS *MELASMETUS* Reitter, 1913: 39

SPECIES *P. femoralis* (Ménétriés, 1832: 228)

Orig. comb.: *Callidium femorale*

Type loc.: "Talyche" (Caucasus)

SUBSPECIES *P. f. demelti* Heyrovsky, 1962: 41

Orig. comb.: *Phymatodes femoralis demelti*

Type loc.: İstanbul (Turkey)

Endemic subspecies to Turkey.

SUBGENUS *PHYMATODELLUS* Reitter, 1913: 40

SPECIES *P. rufipes* (Fabricius, 1777: 232)

Orig. comb.: *Callidium rufipes*

Type loc.: "Germania" (Germany)

SUBSPECIES *P. r. syriacus* (Pic, 1891: 118)

Orig. comb.: *Callidium* (*Poecilium*) *rufipes* var. *syriacum*

Type loc.: "Syrie" (Turkey: ?Hatay: Akbez not Syria)

SUBGENUS *POECILIUM* Fairmaire, 1864: 134

SPECIES *P. kasnaki* (Sama, 2011: 826)

Orig. comb.: *Poecilium kasnaki*

Type loc.: Isparta: Yukarıgökdere (Turkey)

Endemic species to Turkey.

SPECIES *P. magnanii* (Sama & Rapuzzi, 1999: 468)

Orig. comb.: *Poecilium magnanii*

Type loc.: Antalya: Akseki: Çakılı pass (Turkey)

Endemic species to Turkey.

TRIBE ANAGLYPTINI Lacordaire, 1868: 404

SUBTRIBE -

GENUS *ANAGLYPTUS* Mulsant, 1839: 91

SUBGENUS *ANAGLYPTUS* Mulsant, 1839: 91

SPECIES *A. arabicus* (Küster, 1847: 95)

Orig. comb.: *Clytus arabicus*

Type loc.: Turkey

SPECIES *A. croceus* Pesarini & Sabbadini, 1997: 47

Orig. comb.: *Anaglyptus croceus*

Type loc.: İzmir: Bozdağ (Turkey)

Endemic species to Turkey.

TRIBE CLYTINI Mulsant, 1839: 70

SUBTRIBE -

GENUS *ISOTOMUS* Mulsant, 1862: 143

SUBGENUS -

SPECIES *I. comptus* (Mannerheim, 1825: 36)

Orig. comb.: *Clytus comptus*

Type loc.: "Imperii Rossici"

SUBSPECIES *I. c. meridionalis* Özdikmen & Aytar, 2012: 653

Orig. comb.: *Isotomus comptus meridionalis*

Type loc.: Osmaniye: Karatepe (Turkey)

Endemic subspecies to Turkey.

SPECIES *I. syriacus* (Pic, 1902: 25)

Orig. comb.: *Caloclytus syriacus*

Type loc.: "Syrie" (Turkey: Hatay not Syria)

Endemic species to Turkey.

GENUS *CHLOROPHORUS* Chevrolat, 186: 290

SUBGENUS *CRASSOFASCIATUS* Özdikmen, 2011: 538

SPECIES *C. convexifrons* Holzschuh, 1981: 100

Orig. comb.: *Chlorophorus convexifrons*

Type loc.: Samsun and Manisa (Turkey)

Endemic species to Turkey.

SPECIES *C. cursor* Rapuzzi & Sama, 1999: 331

Orig. comb.: *Chlorophorus cursor*

Type loc.: Bolu: Abant Lake (Turkey)

Endemic species to Turkey.

SPECIES *C. niehuisi* Adlbauer, 1992: 497

Orig. comb.: *Chlorophorus niehuisi*

Type loc.: Muş: Buğlan pass (Turkey)

Endemic species to Turkey.

SPECIES *C. oezdikmeni* Sama & Rapuzzi, 2011

Orig. comb.: *Chlorophorus oezdikmeni*

Type loc.: Kahramanmaraş: Andırın (Turkey)

Endemic species to Turkey.

SPECIES *C. robustior* (Pic, 1900: 11)
Orig. comb.: *Clytanthus trifasciatus* var. *robustior*
Type loc.: Konya (Turkey)
Endemic species to Turkey.

SUBGENUS *PERDEROMACULATUS* Özdikmen, 2011: 537

SPECIES *C. graciosus* (Marseul, 1868: 203)
Orig. comb.: *Clytus graciosus*
Type loc.: Beyrouth env. (Lebanon)

SUBSPECIES *C. g. sparsus* (Marseul, 1868: 203)
Orig. comb.: *Clytus (Clytanthus) sparsus*
Type loc.: "Adalia, Lycischen Taurus" (Turkey: Antalya)
Endemic subspecies to Turkey.

SPECIES *C. grosseri* Sama & Rapuzzi, 2011: 85
Orig. comb.: *Chlorophorus grosseri*
Type loc.: Şirnak: Mersindağı pass (Turkey)
Endemic species to Turkey.

SPECIES *C. wewalkai* Holzschuh, 1969: 77
Orig. comb.: *Chlorophorus wewalkai*
Type loc.: Ankara: Kızılcahamam (Turkey)
Endemic species to Turkey.

SUBGENUS *HUMEROMACULATUS* Özdikmen, 2011: 537

SPECIES *C. dinae* Rapuzzi & Sama, 1999: 329
Orig. comb.: *Chlorophorus dinae*
Type loc.: Hatay: Antakya: Şenköy (Turkey)

SPECIES *C. dominici* Sama, 1996: 110
Orig. comb.: *Chlorophorus dominici*
Type loc.: Kastamonu: Devrekani (Turkey)
Endemic species to Turkey.

SPECIES *C. nivipictus* (Kraatz, 1879: 91)
Orig. comb.: *Clytus nivipictus*
Type loc.: "Cilicischen Taurus" (Turkey)

GENUS *TURANOCLYTUS* Sama, 1994: hevrolat, 1860: 456

SUBGENUS -

SPECIES *T. sieversi* (Ganglbauer, 1890: 429)
Orig. comb.: *Clytus (Xylotrechus) sieversi*
Type loc.: "Kasikoparan, (Armenia rossica)" (Turkey: Iğdır)

SYNONYM

akbesianus Pic, 1902: 17
Orig. comb.: *Xylotrechus akbesianus*
Type loc.: Hatay: Akbez (Turkey)

GENUS *PSEUDOSPHEGESTHES* Reitter, 1913: 50

SUBGENUS -

SPECIES *P. longitarsus* Holzschuh, 1974: 90
Orig. comb.: *Pseudosphegesthes longitarsus*
Type loc.: Muş: Buğlan pass (Turkey)
Endemic species to Turkey.

SPECIES *P. samai* Danilevsky, 2000: 44
Orig. comb.: *Pseudosphegesthes samai*
Type loc.: Tokat: Niksar (Turkey)
Endemic species to Turkey.

GENUS *CLYTUS* Laicharting, 1784: 88**SUBGENUS -****SPECIES *C. buglanicus* Kadlec, 2005: 106****Orig. comb.:** *Clytus buglanicus***Type loc.:** Muş; Buğlan pass (Turkey)**Endemic species to Turkey.****SPECIES *C. ciliciensis* (Chevrolat, 1863: 334)****Orig. comb.:** *Sphegesthes ciliciensis***Type loc.:** "Caramanie" (Turkey: Karaman)**SPECIES *C. gulekanus* Pic, 1904: 65****Orig. comb.:** *Clytus gulekanus***Type loc.:** İçel; Gülek (Turkey)**Endemic species to Turkey.****SPECIES *C. insignitus* Fairmaire, 1866: 269 [DA]****Orig. comb.:** *Clytus insignitus***Type loc.:** "Kisilgye-Aole" (Turkey: ?İzmir or Burdur: Kızılkaya)**Endemic species to Turkey.****SPECIES *C. kumalariensis* Johanides, 2001: 219****Orig. comb.:** *Clytus kumalariensis***Type loc.:** Afyon: Kumalar Mt. (Turkey)**Endemic species to Turkey.****SPECIES *C. schneideri* Kiesenwetter, 1878: 313 [= 1879: 57]****Orig. comb.:** *Clytus schneideri***Type loc.:** Caucasus**SUBSPECIES *C. s. inapicalis* Pic, 1895: 38****Orig. comb.:** *Clytus inapicalis***Type loc.:** Erzurum (Turkey)**Endemic subspecies to Turkey.****SPECIES *C. schurmanni* Sama, 1996: 108****Orig. comb.:** *Clytus schurmanni***Type loc.:** Ankara: Kızılcahamam (Turkey)**Endemic species to Turkey.****SPECIES *C. taurusiensis* (Pic, 1903: 139)****Orig. comb.:** *Clytus (Clytanthus) taurusiensis***Type loc.:** "Taurus Mts." (Turkey)**SUBFAMILY STENOPTERINAE Gistel, 1848: [9] (unnumbered section)****TRIBE STENOPTERINI Gistel, 1848: [9]****SUBTRIBE -****GENUS *STENOPTERUS* Illiger, 1804: 120****SUBGENUS -****SPECIES *S. adlbaueri* Sama, 1995: 408****Orig. comb.:** *Stenopterus adlbaueri***Type loc.:** Hakkari (Turkey)**Endemic species to Turkey.****SPECIES *S. atricornis* Pic, 1891: 102****Orig. comb.:** *Stenopterus rufus* v. *atricornis***Type loc.:** "Syrie (Akbès)" (Turkey: Hatay: Akbez not Syria)**SPECIES *S. kraatzi* Pic, 1892: 21****Orig. comb.:** *Stenopterus kraatzi***Type loc.:** "Asie Mineure, Smyrna" (Turkey: İzmir)**Endemic species to Turkey.****SPECIES *S. rufus* (Linnaeus, 1767: 642)****Orig. comb.:** *Necydalis rufus***Type loc.:** "Europa australi" (Southern Europe)

SUBSPECIES *S. r. syriacus* Pic, 1892: 22

Orig. comb.: *Stenopterus rufus* v. *syriacus*

Type loc.: "Syrie (Akbès)" (Turkey: Hatay: Akbez not Syria)

TRIBE MOLORCHINI Gistel, 1848: [9] (unnumbered section)

SUBTRIBE -

GENUS *MOLORCHUS* Fabricius, 1792: 356

SUBGENUS *CAENOPTERA* Thomson, 1859: 150

SPECIES *M. abieticola* Holzschuh, 2007: 218

Orig. comb.: *Molorchus abieticola*

Type loc.: İçel: Namrun (Turkey)

Endemic species to Turkey.

SPECIES *M. juglandis* Sama, 1982: 219

Orig. comb.: *Molorchus juglandis*

Type loc.: Antalya: Alanya (Turkey)

SUBGENUS *MOLORCHUS* Fabricius, 1792: 356

SPECIES *M. kiesenwetteri* Mulsant & Rey, 1861: 189

Orig. comb.: *Molorchus kiesenwetteri*

Type loc.: not stated, probably Germany

SUBSPECIES *M. k. hircus* Abeille de Perrin, 1881: 133

Orig. comb.: *Molorchus hircus*

Type loc.: "Bloudan (Anti Liban)" (Southern Syria)

SYNONYMS

anatolicus Adlbauer, 1988: 275

Orig. comb.: *Molorchus kiesenwetteri anatolicus*

Type loc.: İçel: Erdemli (Turkey)

angorensis Pic, 1912: 3

Orig. comb.: *Caenoptera angorensis*

Type loc.: Ankara (Turkey)

castanipennis Reitter, 1912: 27

Orig. comb.: *Molorchus kiesenwetteri* ab. *castanipennis*

Type loc.: "Kleinasien, Araxestal" (Turkey)

SPECIES *M. malmusii* (Sama, 1995: 370)

Orig. comb.: *Glaphyra malmusii*

Type loc.: Bolu: Abant (Turkey)

Endemic species to Turkey.

SPECIES *M. marmottani* (Brisout, 1863: 118)

Orig. comb.: *Molorchus marmottani*

Type loc.: "Vernet les Bains" (France: Pyrenees)

SUBSPECIES *M. m. frischi* (Sama, 1995: 373)

Orig. comb.: *Glaphyra marmottani frischi*

Type loc.: Osmaniye: Zorkun plateau (Turkey)

Endemic subspecies to Turkey.

SPECIES *M. tenuitarsis* Holzschuh, 1981: 97

Orig. comb.: *Molorchus tenuitarsis*

Type loc.: İçel: Namrun (Turkey)

Endemic species to Turkey.

SYNONYM

sterbai Adlbauer, 1988: 277

Orig. comb.: *Molorchus sterbai*

Type loc.: Kahramanmaraş: Gökşun (Turkey)

TRIBE BRACHYPTEROMINI Sama, 2008: 229**SUBTRIBE -****GENUS** *BRACHYPTEROMA* Heyden, 1863: 128 [NP]**SUBGENUS -****SPECIES** *B. holtzi* Pic, 1905: 114**Orig. comb.:** *Brachypteroma holtzi***Type loc.:** Taurus Mts. (Turkey)**SPECIES** *B. ottomanum* Heyden, 1863: 128 [NP]**Orig. comb.:** *Brachypteroma ottomanum***Type loc.:** European Turkey**TRIBE HYBODERINI** Linsley, 1840: 367**GENUS** *CALLIMUS* Mulsant, 1846: [5]**SUBGENUS** *CALLIMUS* Mulsant, 1846: [5]**SPECIES** *C. akbesianus* Pic, 1892: CXI**Orig. comb.:** *Callimus akbesianus***Type loc.:** Hatay: Akbez (Turkey)**Endemic species to Turkey.****SUBGENUS** *PROCALLIMUS* Pic, 1907: 7**SPECIES** *C. egregius* Mulsant & Rey, 1863: 146**Orig. comb.:** *Callimus egregius***Type loc.:** "La Caramanie" (Turkey: Karaman)

As a result of the present work, a total of 66 species group taxa (51 species and 15 subspecies) of Cerambycidae [Cerambycinae (40 sp. and 13 ssp.) and Stenopterinae (11 sp. and 2 ssp.)] are determined.

Among of them, 44 species group taxa (35 species and 9 subspecies) are endemic to Turkey now. So, endemism ratio of the longhorned beetles that originally described from whole territories of Turkey, is approximately 68 % [Cerambycidae [Cerambycinae (29 sp. and 8 ssp.) and Stenopterinae (6 sp. and 1 ssp.)]].

ACKNOWLEDGEMENT

Special thanks to M. L. Danilevsky (Russia) for some information.

LITERATURE CITED

Löbl I. & Smetana A. (ed.) 2010. Catalogue of Palaearctic Coleoptera, Vol. 6. Chrysomeloidea. Stenstrup: Apollo Books, 924 pp.

Özdikmen, H. 2012a. The longhorned beetles that originally described from whole territories of Turkey (Coleoptera: Cerambycoidea) Part I - Vesperidae and Cerambycidae (Prioninae, Lepturinae, Necydalinae, Aseminae, Saphaninae and Dorcasominae). *Munis Entomology & Zoology*, 7 (1): 592-603.

Özdikmen, H. 2012b. Naked lists of Turkish Cerambycoidea and Chrysomeloidea (Coleoptera). *Munis Entomology & Zoology*, 7 (1): 51-108.